

1257. METHANOCELLA MEDIUM

NaCl	1.00	g
MgCl ₂ x 6 H ₂ O	0.40	g
CaCl ₂ x 2 H ₂ O	0.10	g
NH ₄ Cl	0.10	g
KH ₂ PO ₄	0.20	g
KCl	0.50	g
Trace element solution (see medium 318)	1.00	ml
Yeast extract (OXOID)	0.10	g
Na-acetate	0.10	g
Na-resazurin solution (0.1% w/v)	0.50	ml
NaHCO ₃	2.50	g
Vitamin solution (see medium 141)	10.00	ml
L-Cysteine-HCl x H ₂ O	0.30	g
Na ₂ S x 9 H ₂ O	0.30	g
Distilled water	1000.00	ml

Dissolve ingredients (except bicarbonate, vitamins and reducing agents), then sparge medium with 80% H₂ and 20% CO₂ gas mixture for 30 – 45 min to make it anoxic and adjust pH to 6.0. Dispense medium under same gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. Prior to inoculation add vitamins, cysteine and sulfide from sterile anoxic stock solution prepared under 100% N₂ gas and bicarbonate from a sterile anoxic stock solution prepared under 80% N₂ and 20% CO₂ gas atmosphere. Vitamins are sterilized by filtration. If necessary, adjust the pH of the complete medium to 7.0. After inoculation pressurize vials to 1 bar overpressure with sterile 80% H₂ and 20% CO₂ gas mixture.