

## 1325. PARACOCCLUS BOGORIENSIS MEDIUM

Yeast extract	5.00	g
Peptone	5.00	g
Glucose	2.00	g
KH <sub>2</sub> PO <sub>4</sub>	1.00	g
MgSO <sub>4</sub> x 7 H <sub>2</sub> O	0.20	g
Na <sub>2</sub> CO <sub>3</sub> (7.5% w/v)	40.0	ml
Trace element solution SL10	1.00	ml
Distilled water	960.00	ml

Final pH to 9.5.

Autoclave the Na<sub>2</sub>CO<sub>3</sub> solution separately and combine with the rest of the medium after cooling both to about 50°C. Agar may be added at 15g/l to solidify the medium

### Trace element solution SL10

HCl (concentrated)	5.000	ml
FeCl <sub>2</sub> x 4 H <sub>2</sub> O	1.500	g
ZnCl <sub>2</sub>	0.070	g
MnCl <sub>2</sub> x 4 H <sub>2</sub> O	0.200	g
H <sub>3</sub> BO <sub>3</sub>	0.006	g
CoCl <sub>2</sub> x 6 H <sub>2</sub> O	0.190	g
CuCl <sub>2</sub> x 2 H <sub>2</sub> O	0.002	g
NiCl <sub>2</sub> x 6 H <sub>2</sub> O	0.024	g
Na <sub>2</sub> MoO <sub>4</sub> x 2 H <sub>2</sub> O	0.036	g
Distilled water	1000.000	ml