

14. CLOSTRIDIUM FORMICOACETICUM MEDIUM

Yeast extract	5.00	g
K ₂ HPO ₄	10.00	g
Trace element solution SL-4 (see below)	10.00	ml
Pyridoxine-HCl solution (0.1% w/v)	1.00	ml
Na-thioglycolate	0.75	g
Na-resazurin solution (0.1% w/v)	0.50	ml
NaHCO ₃	10.00	g
D-Fructose	5.00	g
Agar (for solid medium)	15.00	g
Distilled water	1000.00	ml

Dissolve ingredients (except bicarbonate and fructose) and sparge medium with 100% N₂ gas for at least 30 – 45 min to make it anoxic. Dispense medium under same gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. Add fructose from a sterile anoxic stock solution prepared under 100% N₂ gas and bicarbonate from a sterile anoxic stock solution prepared under 80% N₂ and 20% CO₂ gas atmosphere. Adjust pH of complete medium to 8.0.

Trace element solution SL-4:

Na ₂ -EDTA	0.50	g
FeSO ₄ x 7 H ₂ O	0.20	g
ZnSO ₄ x 7 H ₂ O	0.10	g
MnCl ₂ x 4 H ₂ O	0.03	g
H ₃ BO ₃	0.30	g
CoCl ₂ x 6 H ₂ O	0.20	g
CuCl ₂ x 2 H ₂ O	0.01	g
NiCl ₂ x 6 H ₂ O	0.02	g
Na ₂ MoO ₄ x 2 H ₂ O	0.03	g
Distilled water	1000.00	ml

First dissolve EDTA in distilled water and adjust pH to 7.0 using 2 N NaOH; then add other compounds.