

1565: ALKALIPHILUS TRANSVAALENSIS MEDIUM

MJ(-N) synthetic seawater	100.00	ml
Yeast extract (BD Bacto)	2.00	g
Tryptone (BD Bacto)	2.00	g
Na ₂ -tartrate x 2 H ₂ O	1.00	g
Na ₃ -citrate	1.00	g
NaNO ₃	0.25	g
(NH ₄) ₂ SO ₄	0.25	g
Fe(III) citrate	0.20	g
Na ₂ SO ₄	1.00	g
Na ₂ CO ₃	1.25	g
NaHCO ₃	2.50	g
Na ₂ S ₂ O ₃ x 5 H ₂ O	2.00	g
Distilled water	900.00	ml

Dissolve ingredients except carbonate, bicarbonate and thiosulfate, then sparge medium with 100% N₂ gas for 30 - 45 min to make it anoxic. Add and dissolve carbonate and bicarbonate, adjust pH to 11 with KOH, then dispense medium under 100% N₂ gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. Add thiosulfate to the autoclaved medium from an anoxic stock solution prepared under 100% N₂ gas and sterilized by filtration. Adjust pH of complete medium to 9.8 - 10.0.

MJ(-N) synthetic seawater (from medium J268)

NaCl	30.00	g
K ₂ HPO ₄	0.14	g
CaCl ₂ x 2 H ₂ O	0.14	g
MgSO ₄ x 7 H ₂ O	3.40	g
MgCl ₂ x 6 H ₂ O	4.18	g
KCl	0.33	g
NiCl ₂ x 6 H ₂ O	0.50	mg
Na ₂ SeO ₃ x 5 H ₂ O	0.50	mg
Fe(NH ₄) ₂ (SO ₄) ₂ x 6 H ₂ O	0.01	g
Modified Wolin's mineral solution	10.00	ml
Distilled water	1000.00	ml

Modified Wolin's mineral solution (from medium 141)

Nitrolotriacetic acid	1.50	g
MgSO ₄ x 7 H ₂ O	3.00	g
MnSO ₄ x H ₂ O	0.50	g
NaCl	1.00	g
FeSO ₄ x 7 H ₂ O	0.10	g

CoSO ₄ x 7 H ₂ O	0.18	g
CaCl ₂ x 2 H ₂ O	0.10	g
ZnSO ₄ x 7 H ₂ O	0.18	g
CuSO ₄ x 5 H ₂ O	0.01	g
AlK(SO ₄) ₂ x 12 H ₂ O	0.02	g
H ₃ BO ₃	0.01	g
Na ₂ MoO ₄ x 2 H ₂ O	0.01	g
NiCl ₂ x 6 H ₂ O	0.03	g
Na ₂ SeO ₃ x 5 H ₂ O	0.30	mg
Na ₂ WO ₄ x 2 H ₂ O	0.40	mg
Distilled water	1000.00	ml

First dissolve nitrilotriacetic acid and adjust pH to 6.5 with KOH, then add minerals. Adjust final to pH 7.0 with KOH.