

287: METHANOCULLEUS CHIKUGOENSIS MEDIUM

K ₂ HPO ₄	0.30	g
KH ₂ PO ₄	0.30	g
(NH ₄) ₂ SO ₄	0.30	g
NaCl	0.61	g
CaCl ₂ x H ₂ O	0.14	g
MgSO ₄ x 7 H ₂ O	0.13	g
Modified Wolin's mineral solution	10.00	ml
NH ₄ Cl	2.70	g
Na-acetate	2.50	g
Sodium resazurin (0.1% w/v)	0.50	ml
NaHCO ₃	5.00	g
Wolin's vitamin solution (10x)	1.00	ml
L-Cysteine HCl x H ₂ O	0.30	g
Na ₂ S x 9 H ₂ O	0.30	g
Distilled water	1000.00	ml

Dissolve ingredients (except bicarbonate, vitamins, cysteine and sulfide) and sparge medium with 80% H₂ and 20% CO₂ gas mixture for 30 - 45 min to make it anoxic. Add and dissolve bicarbonate, adjust pH to 6.8 and then dispense medium under 80% H₂ and 20% CO₂ gas atmosphere into anoxic Hungate-type tubes or serum bottles and autoclave. After sterilization add cysteine and sulfide from sterile anoxic stock solutions autoclaved under 100% N₂ gas. Vitamins are prepared under 100% N₂ gas atmosphere and sterilized by filtration. Adjust pH of complete medium to 6.8 - 7.0. For incubation use sterile 80% H₂ and 20% CO₂ gas mixture at one atmospheres of pressure.

Modified Wolin's mineral solution (from medium 141)

Nitrolotriacetic acid	1.50	g
MgSO ₄ x 7 H ₂ O	3.00	g
MnSO ₄ x H ₂ O	0.50	g
NaCl	1.00	g
FeSO ₄ x 7 H ₂ O	0.10	g
CoSO ₄ x 7 H ₂ O	0.18	g
CaCl ₂ x 2 H ₂ O	0.10	g
ZnSO ₄ x 7 H ₂ O	0.18	g
CuSO ₄ x 5 H ₂ O	0.01	g
AlK(SO ₄) ₂ x 12 H ₂ O	0.02	g
H ₃ BO ₃	0.01	g
Na ₂ MoO ₄ x 2 H ₂ O	0.01	g
NiCl ₂ x 6 H ₂ O	0.03	g
Na ₂ SeO ₃ x 5 H ₂ O	0.30	mg

Na ₂ WO ₄ x 2 H ₂ O	0.40	mg
Distilled water	1000.00	ml

First dissolve nitrilotriacetic acid and adjust pH to 6.5 with KOH, then add minerals. Adjust final to pH 7.0 with KOH.

Wolin's vitamin solution (10x) (from medium 120)

Biotin	20.00	mg
Folic acid	20.00	mg
Pyridoxine hydrochloride	100.00	mg
Thiamine HCl	50.00	mg
Riboflavin	50.00	mg
Nicotinic acid	50.00	mg
Calcium D-(+)-pantothenate	50.00	mg
Vitamin B ₁₂	1.00	mg
p-Aminobenzoic acid	50.00	mg
(DL)-alpha-Lipoic acid	50.00	mg
Distilled water	1000.00	ml