

## 298. PELOBACTER PROPIONICUS MEDIUM (FRESHWATER)

KH <sub>2</sub> PO <sub>4</sub>	0.20	g
NH <sub>4</sub> Cl	0.25	g
NaCl	1.00	g
MgCl <sub>2</sub> x 6 H <sub>2</sub> O	0.40	g
KCl	0.50	g
CaCl <sub>2</sub> x 2 H <sub>2</sub> O	0.15	g
Trace element solution SL-10 (see medium 320)	1.00	ml
Na-resazurin solution (0.1% w/v)	0.50	ml
NaHCO <sub>3</sub>	2.50	g
2,3-butanediol	0.90	g
Na <sub>2</sub> S x 9 H <sub>2</sub> O	0.36	g
Distilled water	1000.00	ml

Dissolve ingredients except bicarbonate, 2,3-butanediol and sulfide, then sparge medium with 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture for 30 – 45 min to make it anoxic. Dispense medium in anoxic Hungate-type tubes or serum vials under the same gas atmosphere and autoclave. Add 2,3-butanediol and sulfide from sterile anoxic stock solutions prepared under 100% N<sub>2</sub> gas and bicarbonate from a sterile anoxic stock solution prepared under 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture. Prior to use adjust pH of complete medium to 7.2 - 7.5.

*Note: Addition of 10 - 20 mg sodium dithionite per liter (e.g. from 5% (w/v) solution freshly prepared under N<sub>2</sub> and filter-sterilized) may stimulate growth of some strains at the beginning.*

For [DSM 2395](#) replace 2,3-butanediol with 1.00 g/l of polyethylene glycol (molecular weight 106 - 20000).

For [DSM 3246](#) replace 2,3-butanediol with 1.00 g/l acetoin.

For [DSM 6652](#) replace 2,3-butanediol with 2.60 g/l Na-glutarate and add 20.00 ml/l rumen fluid, clarified (see medium 1310) to the autoclaved medium.

For [DSM 6832](#) supplement medium with 10.00 ml/l of a vitamin solution (see medium 141) and replace 2,3-butanediol by 1.00 g/l of quinic acid added from a neutralized anoxic stock solution.

*Continued on next page*

For [DSM 8385](#) supplement medium with 1.00 g/l yeast extract and replace 2,3-butanediol by 2.00 g/l D-galactose added from an anoxic stock solution sterilized by filtration.

For [DSM 8909](#) supplement medium with 1.00 ml/l of a vitamin solution (see medium 503) and replace 2,3-butanediol by 1.40 g/l triethanolamine hydrochloride added from a 1 M stock solution prepared and autoclaved under 100% N<sub>2</sub> gas).

For [DSM 11264](#) supplement medium with 0.20 g/l yeast extract and 10.00 ml/l of a vitamins solution (see medium 141). Replace 2,3-butanediol by 1.24 g/l ethylene glycol added from a separately sterilized stock solution.

For [DSM 17541](#) supplement medium with 10.00 ml/l of a vitamin solution (see medium 141) and 1.00 ml/l of selenite-tungstate solution (see medium 385).

For [DSM 29698](#) replace 2,3-butanediol with 2.20 g/l Na-gluconate.

For [DSM 102354](#) supplement medium with 1.00 ml/l of a vitamins solution (see medium 503) and replace 2,3-butanediol by 0.86 g/l crotonic acid added after autoclaving from an anoxic sterile stock solution neutralized with NaOH.

For [DSM 102355](#) supplement medium with 1.00 ml/l of a vitamins solution (see medium 503) and replace 2,3-butanediol by 1.25 g/l sodium pyruvate added after autoclaving from an anoxic stock solution sterilized by filtration.