

## 298b. SYNTROPHOBOTULUS MEDIUM

KH <sub>2</sub> PO <sub>4</sub>	0.20	g
NH <sub>4</sub> Cl	0.25	g
NaCl	1.00	g
MgCl <sub>2</sub> x 6 H <sub>2</sub> O	0.40	g
KCl	0.50	g
CaCl <sub>2</sub> x 2 H <sub>2</sub> O	0.15	g
Trace element solution SL-10 (see medium 320)	1.00	ml
Na-acetate	0.20	g
Na-resazurin solution (0.1% w/v)	0.50	ml
NaHCO <sub>3</sub>	2.50	g
Na <sub>2</sub> -glyoxalate	0.60	g
Vitamins solution (see medium 141)	10.00	ml
L-Cysteine-HCl x H <sub>2</sub> O	0.30	g
Na <sub>2</sub> S x 9 H <sub>2</sub> O	0.30	g
Distilled water	1000.00	ml

Dissolve ingredients except bicarbonate, glyoxalate, vitamins, cysteine and sulfide, then sparge medium with 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture for 30 – 45 min to make it anoxic. Dispense medium in anoxic Hungate-type tubes or serum vials under the same gas atmosphere and autoclave. Add glyoxalate (sterilized by filtration), vitamins (sterilized by filtration), cysteine and sulfide from sterile anoxic stock solutions prepared under 100% N<sub>2</sub> gas and bicarbonate from a sterile anoxic stock solution prepared under 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture. Prior to use adjust pH of complete medium to 7.2 - 7.5.

*Note: When growth has started, feed culture once or twice with the same amount of glyoxalate.*

For [DSM 6945](#) replace glyoxalate with 1.00 g/l Na-glycolate.