

**311a. SPOROMUSA MEDIUM (H<sub>2</sub>/CO<sub>2</sub>)**

NH <sub>4</sub> Cl	0.50	g
MgSO <sub>4</sub> x 7 H <sub>2</sub> O	0.50	g
CaCl <sub>2</sub> x 2 H <sub>2</sub> O	0.25	g
NaCl	2.25	g
FeSO <sub>4</sub> x 7 H <sub>2</sub> O solution (0.1% w/v in 0.1 N H <sub>2</sub> SO <sub>4</sub> )	2.00	ml
Trace element solution SL-10 (see medium 320)	1.00	ml
Selenite-tungstate solution (see medium 385)	1.00	ml
Yeast extract	2.00	g
Casitone	2.00	g
Na-resazurin solution (0.1% w/v)	0.50	ml
K <sub>2</sub> HPO <sub>4</sub>	0.35	g
KH <sub>2</sub> PO <sub>4</sub>	0.23	g
NaHCO <sub>3</sub>	4.00	g
Vitamin solution (see medium 141)	10.00	ml
DL-Dithiothreitol	0.15	g
Distilled water	1000.00	ml

Dissolve ingredients (except phosphates, bicarbonate, vitamins and dithiothreitol) and sparge medium with 80% H<sub>2</sub> and 20% CO<sub>2</sub> gas mixture for 30 – 45 min to make it anoxic. Dispense under same gas atmosphere in anoxic Hungate-type tubes or serum vials and autoclave. Add phosphates to the medium after autoclaving from sterile stock solutions prepared under 100% N<sub>2</sub> gas and bicarbonate from a sterile stock solution prepared under 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture. Vitamins and dithiothreitol should be prepared under N<sub>2</sub> gas and sterilized by filtration. Adjust pH of completed medium to pH 7.0.

For [DSM 11379](#), [DSM 11380](#), [DSM 11381](#) and [DSM 11382](#) supplement medium with 2.50 g/l Na-pyruvate.