

## 667: DESULFALLAS THERMOSAPOVORANS MEDIUM

<b>Solution A</b>	973.00	ml
<b>Solution B</b>	30.00	ml
<b>Solution C</b>	1.00	ml
<b>Solution D</b>	5.00	ml
<b>Solution E</b>	1.00	ml

Solution A is sparged with 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture to reach a pH below 6 (at least 30 min), then distributed under same gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclaved. Solution B is autoclaved separately under 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas atmosphere. Solutions C is prepared under 100% N<sub>2</sub> gas atmosphere and sterilized by filtration. Solution D is autoclaved under 100% N<sub>2</sub> gas. Solution E is prepared under 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas atmosphere and sterilized by filtration. To complete the medium appropriate amounts of solutions B to E are added to the sterile solution A in the sequence as indicated. Final pH of the medium should be at 7.0 - 7.2.

### Solution A

Na <sub>2</sub> SO <sub>4</sub>	3.00	g
KH <sub>2</sub> PO <sub>4</sub>	0.20	g
NH <sub>4</sub> Cl	1.00	g
NaCl	8.00	g
MgCl <sub>2</sub> x 6 H <sub>2</sub> O	0.40	g
KCl	0.50	g
CaCl <sub>2</sub> x 2 H <sub>2</sub> O	0.15	g
<b>Trace element solution SL-10</b>	1.50	ml
<b>Selenite-tungstate solution</b>	1.00	ml
Na-acetate	0.50	g
Na-butyrate	2.20	g
Yeast extract	0.40	g
Sodium resazurin (0.1% w/v)	0.50	ml
Distilled water	970.00	ml

### Solution B

Na <sub>2</sub> CO <sub>3</sub>	1.50	g
Distilled water	30.00	ml

### Solution C

<b>Wolin's vitamin solution (10x)</b>	1.00	ml
---------------------------------------	------	----

## 667: DESULFALLAS THERMOSAPOVORANS MEDIUM

### Solution D

Na <sub>2</sub> S x 9 H <sub>2</sub> O	0.20	g
Distilled water	5.00	ml

### Solution E

Na-dithionite solution (5% w/v)	1.00	ml
---------------------------------	------	----

### Trace element solution SL-10 (from medium 320)

HCl (25%)	10.00	ml
FeCl <sub>2</sub> x 4 H <sub>2</sub> O	1.50	g
ZnCl <sub>2</sub>	70.00	mg
MnCl <sub>2</sub> x 4 H <sub>2</sub> O	100.00	mg
H <sub>3</sub> BO <sub>3</sub>	6.00	mg
CoCl <sub>2</sub> x 6 H <sub>2</sub> O	190.00	mg
CuCl <sub>2</sub> x 2 H <sub>2</sub> O	2.00	mg
NiCl <sub>2</sub> x 6 H <sub>2</sub> O	24.00	mg
Na <sub>2</sub> MoO <sub>4</sub> x 2 H <sub>2</sub> O	36.00	mg
Distilled water	990.00	ml

First dissolve FeCl<sub>2</sub> in the HCl, then dilute in water, add and dissolve the other salts. Finally make up to 1000.00 ml.

### Selenite-tungstate solution (from medium 385)

NaOH	0.50	g
Na <sub>2</sub> SeO <sub>3</sub> x 5 H <sub>2</sub> O	3.00	mg
Na <sub>2</sub> WO <sub>4</sub> x 2 H <sub>2</sub> O	4.00	mg
Distilled water	1000.00	ml

### Wolin's vitamin solution (10x) (from medium 120)

Biotin	20.00	mg
Folic acid	20.00	mg
Pyridoxine hydrochloride	100.00	mg
Thiamine HCl	50.00	mg
Riboflavin	50.00	mg
Nicotinic acid	50.00	mg
Calcium D-(+)-pantothenate	50.00	mg
Vitamin B <sub>12</sub>	1.00	mg
p-Aminobenzoic acid	50.00	mg
(DL)-alpha-Lipoic acid	50.00	mg
Distilled water	1000.00	ml

### Na-dithionite solution (5% w/v) (from medium 829)

NaHCO <sub>3</sub>	50.00	g
Na <sub>2</sub> S <sub>2</sub> O <sub>4</sub>	50.00	g
Distilled water	1000.00	ml

Dissolve NaHCO<sub>3</sub> in water and make the solution anoxic by sparging with 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture. Then dissolve the Na-dithionite and filter sterilize the solution into anoxic Hungate tubes. Store the prepared solution in the dark and refrigerated. Prepare only small amounts of stock solution, as Na-dithionite decomposes rapidly.