

69. THIOBACILLUS NOVELLUS MEDIUM

Solution A:

Na ₂ HPO ₄ x 12 H ₂ O	10.60	g
KH ₂ PO ₄	1.50	g
NH ₄ Cl	0.30	g
Yeast extract	0.30	g
Phenol red	2.00	mg
Distilled water	900.00	ml

Solution B:

MgSO ₄ x 7 H ₂ O	0.10	g
Distilled water	50.00	ml

Solution C:

Trace element solution (see below)	5.00	ml
------------------------------------	------	----

Solution D:

Na ₂ S ₂ O ₃ x 5 H ₂ O	5.00	g
Distilled water	50.00	ml

Solutions A, B, C, D are sterilized separately by autoclaving and mixed when cold. For agar medium add 1.5% agar (Difco Bacto) to solution A. Adjust pH to 8.5 with sterile 0.5 N NaOH.

Trace element solution (Vishniac and Santer, 1957):

Na ₂ -EDTA	50.00	g
ZnSO ₄ x 7 H ₂ O	22.00	g
CaCl ₂ x 2 H ₂ O	5.54	g
MnCl ₂ x 4 H ₂ O	5.06	g
FeSO ₄ x 7 H ₂ O	5.00	g
(NH ₄) ₆ Mo ₇ O ₂₄ x 4 H ₂ O	1.10	g
CuSO ₄ x 5 H ₂ O	1.57	g
CoCl ₂ x 6 H ₂ O	1.61	g
Distilled water	1000.00	ml

Adjust pH to 6.0 with KOH.