

**813a. DESULFONATRONUM THIODISMUTANS MEDIUM**

KH <sub>2</sub> PO <sub>4</sub>	0.2	g
MgCl <sub>2</sub> x 6 H <sub>2</sub> O	0.1	g
NH <sub>4</sub> Cl	1.0	g
KCl	0.2	g
Na <sub>2</sub> SO <sub>4</sub>	5.0	g
NaCl	30.0	g
Trace element solution SL-10 (see medium 320)	1.0	ml
Yeast extract	1.0	g
Na-resazurin solution (0.1% w/v)	0.5	ml
Na <sub>2</sub> CO <sub>3</sub>	3.5	g
Na <sub>2</sub> S x 9 H <sub>2</sub> O	0.5	g
Vitamin solution (see medium 141)	10.0	ml
Na-formate	4.0	g
Distilled water	1000.0	ml

Dissolve the ingredients (except carbonate, sulfide, vitamins and formate) and sparge medium with 100% N<sub>2</sub> gas for 30 – 45 min to make it anoxic. Add and dissolve carbonate and sulfide, then adjust pH to 9.0 - 9.5, dispense under 100% N<sub>2</sub> gas atmosphere into Hungate-type tubes or serum vials and autoclave. Before use add vitamins (sterilized by filtration) and formate from sterile anoxic stock solutions prepared under 100% N<sub>2</sub> gas.