

## 872: PAPILLIBACTER MEDIUM

NH <sub>4</sub> Cl	0.30	g
K <sub>2</sub> HPO <sub>4</sub>	0.20	g
MgCl <sub>2</sub> x 6 H <sub>2</sub> O	0.40	g
CaCl <sub>2</sub> x 2 H <sub>2</sub> O	0.10	g
KCl	0.50	g
NaCl	1.00	g
Yeast extract (BD Bacto)	5.00	g
Trypticase peptone (BD BBL)	5.00	g
<b>Trace element solution SL-10</b>	1.00	ml
Sodium resazurin (0.1% w/v)	0.50	ml
Na <sub>2</sub> CO <sub>3</sub>	2.00	g
trans-Cinnamic acid	0.75	g
L-Cysteine HCl x H <sub>2</sub> O	0.50	g
Na <sub>2</sub> S x 9 H <sub>2</sub> O	0.50	g
Distilled water	1000.00	ml

Dissolve all ingredients except carbonate, cinnamic acid, sulfide and cysteine, then sparge medium with 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture for 30 - 45 min to make it anoxic. Dispense medium under same gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. Add cinnamic acid, sulfide and cysteine from sterile anoxic stock solutions prepared under 100% N<sub>2</sub> gas and carbonate from a sterile anoxic stock solution prepared under 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas atmosphere. Adjust pH of complete medium to 7.2 - 7.5, if necessary.

### Trace element solution SL-10 (from medium 320)

HCl (25%)	10.00	ml
FeCl <sub>2</sub> x 4 H <sub>2</sub> O	1.50	g
ZnCl <sub>2</sub>	70.00	mg
MnCl <sub>2</sub> x 4 H <sub>2</sub> O	100.00	mg
H <sub>3</sub> BO <sub>3</sub>	6.00	mg
CoCl <sub>2</sub> x 6 H <sub>2</sub> O	190.00	mg
CuCl <sub>2</sub> x 2 H <sub>2</sub> O	2.00	mg
NiCl <sub>2</sub> x 6 H <sub>2</sub> O	24.00	mg
Na <sub>2</sub> MoO <sub>4</sub> x 2 H <sub>2</sub> O	36.00	mg
Distilled water	990.00	ml

First dissolve FeCl<sub>2</sub> in the HCl, then dilute in water, add and dissolve the other salts. Finally make up to 1000.00 ml.