

**931. ALCALIPHILIC AMPHIBACILLUS STRAINS MEDIUM**

KH <sub>2</sub> PO <sub>4</sub>	0.200	g
MgCl <sub>2</sub>	0.100	g
NH <sub>4</sub> Cl	0.500	g
KCl	0.200	g
Na <sub>2</sub> CO <sub>3</sub>	63.600	g
NaHCO <sub>3</sub>	50.400	g
Na <sub>2</sub> S x 9 H <sub>2</sub> O	0.700	g
Yeast extract	0.200	g
Sucrose	5.000	g
Trace elements (see DSM medium 141)	1.000	ml
Vitamin solution (see DSM medium 141)	10.000	ml
Resazurin	0.001	g
Distilled water	1000.000	ml

Final pH 9.5-10.0

Prepare the medium without adding the vitamins, sucrose, yeast extract, NH<sub>4</sub>Cl, Na<sub>2</sub>CO<sub>3</sub>, and NaHCO<sub>3</sub> using anaerobic conditions, under nitrogen. If the medium has been boiled to remove oxygen add the NH<sub>4</sub>Cl, Na<sub>2</sub>CO<sub>3</sub>, and NaHCO<sub>3</sub> after the medium had cooled. Dispense into tubes stopper with rubber stoppers (serum tubes or bottles, or screw capped tubes). Autoclave, and to the cooled medium add the vitamins (filter sterilised under N<sub>2</sub>), sucrose, yeast extract, and Na<sub>2</sub>S x 9 H<sub>2</sub>O from anaerobic, sterile stock solutions. DSM 13869 and DSM 13870 were deposited as anaerobic cultures.