

144. THERMOANAEROBACTER BROCKII MEDIUM

NH ₄ Cl	0.90	g
NaCl	0.90	g
MgCl ₂ x 6 H ₂ O	0.40	g
KH ₂ PO ₄	0.75	g
K ₂ HPO ₄	1.50	g
Trace element solution (see below)	9.00	ml
FeSO ₄ x 7 H ₂ O solution (0.1% w/v in 0.1 N H ₂ SO ₄)	3.00	ml
Yeast extract	3.00	g
Trypticase peptone (BD BBL)	10.00	g
Na-resazurin solution (0.1% w/v)	0.50	ml
Vitamin solution (see medium 141)	5.00	ml
D-Glucose	5.00	g
Na ₂ S x 9 H ₂ O	1.00	g
Distilled water	1000.00	ml

Dissolve ingredients (except vitamins, glucose and sulfide), sparge medium with 100% N₂ gas for 30 – 45 min to make it anoxic. Dispense medium under same gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. Add glucose, vitamins (sterilized by filtration) and sulfide from sterile anoxic stock solutions prepared under 100% N₂ gas. Adjust pH of complete medium to 7.2 - 7.4.

Trace element solution:

Nitrilotriacetic acid (NTA)	12.80	g
FeCl ₂ x 4 H ₂ O	0.20	g
MnCl ₂ x 4 H ₂ O	0.10	g
CoCl ₂ x 6 H ₂ O	0.17	g
CaCl ₂ x 2 H ₂ O	0.10	g
ZnCl ₂	0.10	g
CuCl ₂	0.02	g
H ₃ BO ₃	0.01	g
Na ₂ MoO ₄ x 2 H ₂ O	0.01	g
NiCl ₂ x 6 H ₂ O	0.03	g
NaCl	1.00	g
Na ₂ SeO ₃ x 5 H ₂ O	0.03	g
Distilled water	1000.00	ml

First dissolve NTA in 200 ml of distilled water and adjust pH to 6.5 with KOH, then dissolve mineral salts. Finally adjust pH to 6.5 with KOH and make up to 1000.00 ml.

For [DSM 12299](#) omit D-glucose.