

## 1459. THERMOHALOBACTER MEDIUM

K <sub>2</sub> HPO <sub>4</sub>	0.30	g
KH <sub>2</sub> PO <sub>4</sub>	0.30	g
NH <sub>4</sub> Cl	0.10	g
MgCl <sub>2</sub> x 6 H <sub>2</sub> O	2.00	g
CaCl <sub>2</sub> x 2 H <sub>2</sub> O	0.20	g
NaCl	100.00	g
KCl	4.00	g
Na-acetate	0.50	g
Yeast extract (OXOID)	1.00	g
Trypticase peptone (BD BBL)	1.00	g
Wolfe's mineral elixir (see medium 792)	1.00	ml
Na-resazurin solution (0.1% w/v)	0.50	ml
NaHCO <sub>3</sub>	2.50	g
D-Glucose	10.00	g
L-Cysteine-HCl x H <sub>2</sub> O	0.50	g
Na <sub>2</sub> S x 9 H <sub>2</sub> O	0.50	g
Distilled water	1000.00	ml

Dissolve ingredients except bicarbonate, glucose, cysteine and sulfide, then sparge medium with 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture for 30 – 45 min to make it anoxic. Dispense medium under same gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. Add glucose, cysteine and sulfide from sterile anoxic stock solutions prepared under 100% N<sub>2</sub> gas and bicarbonate from a sterile anoxic stock solution prepared under 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture. Prior to use adjust pH of complete medium to 7.0 - 7.2, if necessary.