

330: RUMEN BACTERIA MEDIUM

Mineral solution	38.00	ml
K ₂ HPO ₄	0.30	g
Trypticase peptone (BD BBL)	2.00	g
Yeast extract (OXOID)	0.50	g
Volatile fatty acid mixture	3.10	ml
Haemin solution (0.05% w/v)	2.00	ml
Glycerol	0.50	g
Sodium resazurin (0.1% w/v)	0.50	ml
Na ₂ CO ₃	4.00	g
D-Glucose	0.50	g
Maltose	0.50	g
Cellobiose	0.50	g
Starch, soluble	0.50	g
L-Cysteine HCl x H ₂ O	0.25	g
Na ₂ S x 9 H ₂ O	0.25	g
Distilled water	960.00	ml

Dissolve ingredients (except carbonate, glucose, maltose, cellobiose, soluble starch, cysteine and sulfide), then sparge medium with 100% CO₂ gas for 30 - 45 min to make it anoxic. Add the carbonate and equilibrate the medium with the CO₂ gas to pH 6.8. Distribute under 100% CO₂ gas atmosphere into anoxic Hungate-type tubes or serum vials and autoclave. Thereafter, add glucose, maltose, cellobiose, soluble starch, cysteine and sulfide from sterile anoxic stock solutions prepared under 100% N₂ gas atmosphere. Adjust pH of complete medium to 6.7 - 6.8, if necessary.

For DSM 17630: Plus rumen fluid (30ml/l)

Mineral solution (from medium 330)

KH ₂ PO ₄	6.00	g
NaCl	12.00	g
(NH ₄) ₂ SO ₄	6.00	g
CaCl ₂ x 2 H ₂ O	1.60	g
MgSO ₄ x 7 H ₂ O	2.50	g
Distilled water	1000.00	ml

Volatile fatty acid mixture (from medium 330)

Acetic acid	548.50	ml
Propionic acid	193.50	ml
Butyric acid	129.00	ml
n-Valeric acid	32.25	ml
iso-Butyric acid	32.25	ml

330: RUMEN BACTERIA MEDIUM

DL-2-Methylbutyric acid	32.25	ml
iso-Valeric acid	32.25	ml

Haemin solution (from medium 78)

Haemin	50.00	mg
NaOH (1 N)	1.00	ml
Distilled water	100.00	ml

Dissolve 50 mg haemin in 1 ml 1 N NaOH; make up to 100 ml with distilled water and filter sterilize. Store refrigerated.