

**647a. DESULFUROMONAS ACETEXIGENES MEDIUM**

NaCl	20.0	g
KH <sub>2</sub> PO <sub>4</sub>	1.0	g
NH <sub>4</sub> Cl	0.5	g
MgSO <sub>4</sub> x 7 H <sub>2</sub> O	1.0	g
MgCl <sub>2</sub> x 6 H <sub>2</sub> O	2.0	g
CaCl <sub>2</sub> x 2 H <sub>2</sub> O	0.1	g
Trace element solution SL-10 (see medium 320)	1.0	ml
Selenite-tungstate solution (see medium 385)	1.0	ml
Na-resazurin solution (0.1% w/v)	0.5	ml
Sulfur, powdered	5.0	g
Na <sub>2</sub> CO <sub>3</sub>	1.5	g
Na-pyruvate	0.6	g
Vitamin solution (see medium 503)	1.0	ml
Na <sub>2</sub> S x 9 H <sub>2</sub> O	0.5	g
Distilled water	1000.0	ml

Dissolve ingredients (except sulfur, carbonate, pyruvate, vitamins and sulfide), then sparge medium with 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas mixture for 30 – 45 min to make it anoxic. Dispense medium under same gas atmosphere into anoxic Hungate-type tubes or serum vials that contain already the appropriate amount of sulfur. Sterilize medium by heating cultivation vessels in a boiling water bath for 2 - 3 hours on each of 3 successive days. Complete the medium by adding pyruvate, vitamins and sulfide from sterile anoxic stock solutions prepared under 100% N<sub>2</sub> gas atmosphere and carbonate from a sterile anoxic stock solution prepared under 80% N<sub>2</sub> and 20% CO<sub>2</sub> gas atmosphere. Pyruvate and vitamins are sterilized by filtration. Adjust pH of the complete medium to 7.0 - 7.2, if necessary.